

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

with the Engelmann herbarium. This unpublished and undistributed material, containing about 650 numbers, represents the collections of 1849–1851, and proves to be very valuable. It has now been published by Blankinship,³³ who has still further added to the value of the contribution by including also the numbers of the earlier fascicles not previously enumerated (*Plantae Lindheimerianae* having been left unfinished at the end of the Compositae); a bibliography of Texan botany; a complete index of all three parts, with modern equivalents and corrections, the nomenclature comforming to the Vienna code; and a most interesting sketch, with portrait, of "Lindheimer, the botanist-editor," from data largely supplied by his son and daughter.—J. M. C.

Mutation and geographic distribution.—Willis³⁴ has continued his arguments in favor of mutation by analyzing the geographic distribution of the Dilleniaceae, stating that this family is chosen simply because it is the first family in Hooker's *Flora of British India* "with other than world-wide distribution." The details of the analysis cannot be given here, but the results are intended to show that the theory of mutation greatly simplifies the problems of geographic distribution.

In another short paper³⁵ Willis suggests what seems to be an important consideration in the origin of species of flowering plants, namely, that "while the characters that distinguish species and genera are largely characters of the floral organs, the struggle for existence is almost entirely among the seedlings and young plants, in which these organs are not yet present."—J. M. C.

Fertilization in Polytrichum.—The VAN LEEUWEN-REYNVAANS³⁶ have published the first account of the details of fertilization in mosses and describe most remarkable behavior by the chromatin. In the next to the last division of the spermatogenous cells each daughter nucleus receives six chromosomes, but in the final mitosis only three, so that the sperm contains only three chromosomes. The mitosis which forms the egg and ventral canal cell shows only three chromosomes for each nucleus. The egg and ventral canal cell become pressed together and their nuclei fuse, forming a nucleus with six chromosomes. Two sperms then unite with this egg, thus restoring the sporophytic number of chromosomes, which was found to be twelve. The full paper with the plates will be awaited with interest.—Charles J. Chamberlain.

³³ Blankinship, J. W., Plantae Lindheimerianae, Part III. Ann. Rep. Mo. Bot. Garden 18:123-223. 1907.

³⁴ WILLIS, J. C., The geographical distribution of the Dilleniaceae, as illustrating the treatment of this subject on the theory of mutation. Annals Bot. Gard. Peradeniya 4:69–76. 1907.

 $^{^{35}\,\}mathrm{Further}$ evidence against the origin of species by infinitesimal variations. Idem $_{17-19}.$

³⁶ VAN LEEUWEN-REYNVAAN, Mr. and Mrs. Doctors, On a double reduction of the number of chromosomes during the formation of the sexual cells and on a subsequent double fertilization in some species of Polytrichum. Koninklijke Akad. Wetenschappen 1907:359–365.